

SUGGESTIONS FOR MANAGEMENT OF ANTICONVULSANT-ANTIRETROVIRAL INTERACTIONS IN HIV

Avoid: carbamazepine, phenytoin, phenobarbital, primidone, felbamate, oxcarbazepine

- **all are enzyme inducers and can decrease protease inhibitor and NNRTI levels, which may impact viral efficacy.**
- In the U.S. Military HIV Natural History Study, virologic response was assessed in 19 patients taking enzyme-inducing (EI) antiepileptics (AEDs) (n=12 phenytoin, n=6 carbamazepine, n=1 phenobarbital) versus 85 patients taking non-enzyme inducing antiepileptics (n=88 gabapentin, n=2 pregabalin, n=1 levetiracetam) while on HAART concomitantly for at least 28 consecutive days. Virologic failure was defined as having 2 consecutive viral loads ≥ 400 copies/mL after 6 months of HAART or having all VLs in the first 6 months of HAART ≥ 400 c/mL. Patients on EI-antiepileptics had significantly greater virologic failure (10/16, 63%) compared to other AED patients (20/75, 27%) for the first HAART/AED period (OR 4.6 [1.5-14.3]; $P < 0.01$). Average VL was also greater during HAART in the EI-AED group (3.3 log \pm 1.3) than the other AED group (2.4 log \pm 1.2; $P < 0.01$). A multivariate model adjusting for both year starting and VL at HAART showed similar results (OR 4.7 [0.9-23.6]; $P = 0.06$). Analysis of multiple overlap periods yielded consistent and significant results with higher rates of VF in the EI-AED group (OR 4.2 [1.5-11.4]; $P < 0.01$). Therefore, concurrent use of enzyme-inducing anti-epileptics and HAART should be avoided whenever possible.[Okulicz et al. 2011]

- 1) Depending on seizure type, **consider using other 2nd-line anticonvulsants** to minimize interactions with protease inhibitors and NNRTIs.

Best choices:

- Gabapentin
- Lamotrigine- a significant decrease in lamotrigine concentrations may be seen when used with ritonavir-based regimens
- Levetiracetam
- Vigabatrin
- Pregabalin

Caution Warranted:

- Valproic acid- monitor viral load closely; avoid combination with zidovudine due to cases of severe anemia
- Zonisamide- potential for increased zonisamide concentrations; a decreased dose may be required
- Topiramate- small potential for increased topiramate concentrations; topiramate is a mild CYP3A4 inducer (may impact PI and NNRTI concentrations)
- Tiagabine- potential for increased tiagabine concentrations; a decreased dose may be required
- Anticonvulsants and protease inhibitors- potential for additive bone toxicity (osteonecrosis, osteopenia)

- 2) **Change antiviral or drug dose if possible:**

- If possible, consider using a raltegravir-based regimen.
- Use ritonavir boosted protease inhibitor regimens (minimum ritonavir) 200mg/day to overcome induction. This is still a preliminary recommendation, since data are limited with this approach. Therapeutic drug monitoring of antiretrovirals is recommended if available.

- Empirically increase Kaletra (lopinavir/ritonavir) dose to 3 tablets BID (i.e. 600/150 mg BID) when combined with enzyme inducing anticonvulsants. This is still a preliminary recommendation, since data are limited with this approach. Therapeutic drug monitoring of both the antiretrovirals and anticonvulsants is recommended if available (see table for details).
- Preliminary evidence suggest that darunavir/ritonavir (DRV/RTV) combined with carbamazepine may be a viable option. Although it is unlikely the dose of DRV/RTV requires adjustments, a 25-50% decrease in carbamazepine dose may be required. Therapeutic drug monitoring of both the antiretrovirals and carbamazepine is recommended if available (see table for details).

Appendix I: Antiepileptic Drugs of Choice in the General Population

Seizure Type	First-line Therapy	Second-line Therapy
<i>a) Generalized</i>		
Tonic-Clonic	VPA, CBZ, PHT	CLB, FBM, ?GBP, LTG, ?LEV, OXC, PB, PRM, ?TGB, TPM, ?VGB, ZNS
Absence	VPA, ESM	AZM, CLB, ?FBM, LTG, ?LEV, ?TPM, ZNS
Myoclonic	VPA	AZM, CLB, CZP, ?FBM, ?LTG, ?LEV, ?TPM, ZNS
Atonic	VPA	CLB, FBM, LTG, TGM
<i>b) Partial</i>		
Simple or complex ± secondary generalization	CBZ, PHT	CLB, FBM, GBP, LTG, LEV, OXC, PB, PGB, PRM, TGB, TPM, VGB, VPA, ZNS
Benign rolandic epilepsy	CBZ	CLB, GBP, PHT
Juvenile myoclonic epilepsy	VPA	CLB, LTG, TPM, CBZ, PHT (for gen. T-C seizures)
Lennox-Gastaut	VPA	CLB, CZP, FBM, LTG, TPM, VGB
West Syndrome (infantile spasms)	VGB superior toACTH	CLB, CZP, LTG, steroids, VPA
<i>c) Women</i>		
Idiopathic generalized epilepsy	VPA	
Idiopathic generalized epilepsy (pregnancy)	LTG	

ACTH= adrenocorticotrophic hormone; AZM= acetazolamide; CBZ= carbamazepine; CLB= clobazam; CZP= clonazepam; ESM= ethosuximide; FBM= felbamate; GBP= gabapentin; LTG= lamotrigine; LEV= levetiracetam; OXC= oxcarbazepine; PB= phenobarbital; PGB= pregabalin; PHT= pheytoin; PRM= primidone; TGB= tiagabine; TPM= topiramate; VGB= vigabatrin; VPA= valproic acid; ZNS= zonisamide. ? = unclear

References:

Guberman A, Bruni J. Documented acceptable monotherapy of newer antiepileptics in new onset and partial refractory epilepsy: LTG, OXC, TPM Essentials of Clinical Epilepsy 2nd Ed. Butterworth-Heinemann, Woburn, MA, 1999.

French JA, Kanner AM, Bautista J, Abou-Khalil B, Brown T, Harden CL, Theodore WH, et al. Efficacy and tolerability of the new antiepileptic drug, Prepared by: Michelle Foisy, Pharm.D., Northern Alberta Program, Royal Alexandra Hospital, Edmonton, Alberta. Updated by Michelle Foisy, Pharm.D., and Alice Tseng, Pharm.D., Toronto General Hospital, April 2011 www.hivclinic.ca Page 2 of 3

I: treatment of new-onset epilepsy: report of the TTA and QSS subcommittees of the American Academy of Neurology and the American Epilepsy Society. *Epilepsia* 2004;45(5):401-409.

French JA, Kanner AM, Bautista J, Abou-Khalil B, Brown T, Harden CL, Theodore WH, et al. Efficacy and tolerability of the new antiepileptic drug, II: treatment of refractory epilepsy: report of the TTA and QSS subcommittees of the American Academy of Neurology and the American Epilepsy Society. *Epilepsia* 2004;45(5):410-423.

Lietke MD, Lockhart SM, Rathbun RC. Anticonvulsant and antiretroviral interactions. *Ann Pharmacother* 2004;38:482-9.

Okulicz J, Grandits G, French J, Simpson D, George J, Weintrob A, et al. Co-administered HAART and CYP450 EI-AED: implications for HIV/epilepsy treatment in resource-limited settings [abstract 646]. 18th Conference on Retroviruses and Opportunistic Infections. Feb 27-Mar 2, 2011, Boston, USA.

Romanelli F, Pomeroy C. Concurrent use of antiretrovirals and anticonvulsants in human immunodeficiency virus (HIV) seropositive patients. *Curr Pharm Des* 2003;9:1433-9.

Tomson T. Drug selection for the newly diagnosed patient: when is a new generation antiepileptic drug indicated? *J Neurol* 2004;251:1043-1049.